

## Original Article

# Prevalence of *bla*<sub>CTX-M</sub> Gene in Multi-Resistant *Escherichia coli* Isolated from Urinary Tract Infections, Tehran, Iran

Mehdi Goudarzi<sup>1,2</sup>, Fattaneh Sabzehali<sup>2</sup>, Zahra Tayebi<sup>3</sup>, Mehdi Azad<sup>4</sup>, Shahram Boromandi<sup>5</sup>, Ali Hashemi<sup>2</sup>, Sima Sadat Seyedjavadi<sup>6\*</sup>

<sup>1</sup> Infectious Diseases and Tropical Medicine Research Center, Shahid Beheshti University of Medical Science, Tehran, Iran

<sup>2</sup> Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Science, Tehran, Iran

<sup>3</sup> Department of Microbiology, Tehran Medical Branch, Islamic Azad University, Tehran, Iran

<sup>4</sup> Faculty of paramedical sciences, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>5</sup> Department of Microbiology, Pars hospital laboratory

<sup>6</sup> Department of Pharmaceutical Biotechnology, Pasteur institute, Tehran, Iran

## Abstract

**Background:** The emergence and increase in the incidence of Extended-spectrum beta lactamase (ESBL) producing *Escherichia coli* (*E. coli*) has become an emerging challenge especially in hospitalized patients with urinary tract infection (UTI). The aim of the present study was to survey the frequency of *bla* CTX-M genotype in ESBL producing *E. coli* isolated from hospitalized patients with urinary tract infection and determination of their antibiotic resistance pattern.

**Materials and Methods:** A total of 135 *E. coli* isolates were collected and isolated from patients with UTI. The isolates were subjected to confirmatory phenotype tests for the presence of ESBL. 75 *E. coli* isolates were confirmed as ESBL-positive by double disc synergy test. In vitro susceptibility of ESBL isolates to 15 antimicrobial agents amoxicillin, penicillin, ceftazidime, cefotaxime, cefoxitin, ceftriaxone, cefixime, cephalixin, co-trimoxazole, gentamicin, nalidixic acid, ciprofloxacin, nitrofurantoin, amikacin, and imipenem was performed by Kirby-Bauer's Disk diffusion method according to Clinical and Laboratory Standards Institute (CLSI, 2012) guideline. PCR method was used to identify *bla* CTX-M gene in 75 ESBL positive strains.

**Results:** PCR and sequence analysis showed that 75 (55.5%) isolates produced *bla* CTX-M genes. In vitro susceptibility of ESBL producing *E. coli* showed that all of them were resistant to amoxicillin and penicillin. The rates of resistance to the majority of tested antibiotics varied among 61% to 100 %, with the exception of amikacin (14.7%) and imipenem (2.7%). Our results showed that the frequency of *bla* CTX-M was strikingly high (93.3%) in patients with UTI.

**Conclusion:** These data confirmed that the frequency of *bla* CTX-M genes was high among *E. coli* isolated from patients with UTI. The trend of multidrug-resistant profile has been associated with *bla* CTX-M gene is alarming. Therefore, it is very important to establish a routine screening of ESBL in clinical isolates to prevent dissemination of resistant isolates in health care settings.

**Keywords:** ESBL, Beta lactamase, *Escherichia coli*, antimicrobial resistance

\*Corresponding Author: Sima Sadat Seyedjavadi, Department of Pharmaceutical Biotechnology, Pasteur institute, Tehran, Iran. Email: Sima\_seyedjavadi@yahoo.com

**Please cite this article as:** Goudarzi M, Sabzehali F, Tayebi Z, Azad M, Boromani SH, Hashemi A, Seyedjavadi SS. Prevalence of *bla*<sub>CTX-M</sub> Gene in Multi-Resistant *Escherichia coli* Isolated from Urinary Tract Infections, Tehran, Iran. Novel Biomed. 2014;2(4):107-13.